

# Homology of the hypertree poset

Bérénice Oger

Institut Camille Jordan (Lyon)

February, 11th 2015

## Motivation : $P\Sigma_n$

- $F_n$  generated by  $(x_i)_{i=1}^n$
- $P\Sigma_n$ , pure symmetric automorphism group
  - ▶ group of automorphisms of  $F_n$  which send each  $x_i$  to a conjugate of itself,
  - ▶ group of motions of a collection of  $n$  coloured unknotted, unlinked circles in 3-space.
- Their cohomology groups are not Koszul (A. Conner and P. Goetz).

- Action of  $P\Sigma_n$  on a contractible complex  $MM_n$  defined by McCullough and Miller in 1996 in terms of marking of hypertrees, whose fundamental domain is the hypertree poset on  $n$  vertices,
- Use of the **hypertree poset** for the computation of the  $l^2$ -Betti numbers of  $P\Sigma_n$  by C. Jensen, J. McCammond and J. Meier,
- $P\Sigma_n \triangleright \text{Inn}(F_n) \Rightarrow OP\Sigma_n = P\Sigma_n / \text{Inn}(F_n)$
- $OP\Sigma_n$  acts cocompactly on  $MM_n$
- Use of a theorem by Davis, Januszkiewicz and Leary to obtain the expression of  $l^2$ -cohomology of the group in term of the cohomology of the fundamental domain of the complex.

# Summary

- 1 The hypertree poset
  - Hypertrees
  - Hypertree poset
  - Homology of the hypertree poset
- 2 Computation of the homology of the hypertree poset
  - Species
  - Counting strict chains using large chains
  - Pointed hypertrees
  - Relations between chains of hypertrees
  - Dimension of the homology
- 3 From the hypertree poset to rooted trees
  - PreLie species
  - Character for the action of the symmetric group on the homology of the poset


# Hypergraphs and hypertrees

## Definition ([Ber89])

A *hypergraph* (on a set  $V$ ) is an ordered pair  $(V, E)$  where:

- $V$  is a finite set (*vertices*)
- $E$  is a collection of subsets of cardinality at least two of elements of  $V$  (*edges*).

## Example of a hypergraph on $[1; 7]$



# Walk on a hypergraph

## Definition

Let  $H = (V, E)$  be a hypergraph.


A *walk* from a vertex or an edge  $d$  to a vertex or an edge  $f$  in  $H$  is an alternating sequence of vertices and edges beginning by  $d$  and ending by  $f$ :

$$(d, \dots, e_i, v_i, e_{i+1}, \dots, f)$$

where for all  $i$ ,  $v_i \in V$ ,  $e_i \in E$  and  $\{v_i, v_{i+1}\} \subseteq e_i$ .

The *length* of a walk is the number of edges and vertices in the walk.

## Examples of walks




# Hypertrees

## Definition

A *hypertree* is a non-empty hypergraph  $H$  such that, given any distinct vertices  $v$  and  $w$  in  $H$ ,

- there exists a walk from  $v$  to  $w$  in  $H$  with distinct edges  $e_i$ , ( $H$  is *connected*),
- and this walk is unique, ( $H$  has *no cycles*).

## Example of a hypertree



# The hypertree poset


## Definition

Let  $I$  be a finite set of cardinality  $n$ ,  $S$  and  $T$  be two hypertrees on  $I$ .


$S \preceq T \iff$  Each edge of  $S$  is the union of edges of  $T$

We write  $S \prec T$  if  $S \preceq T$  but  $S \neq T$ .

## Example with hypertrees on four vertices







- Graded poset by the number of edges [McCullough and Miller 1996],
- There is a unique minimum  $\hat{0}$ ,
- $\text{HT}(I)$  = hypertree poset on  $I$ ,
- $\text{HT}_n$  = hypertree poset on  $n$  vertices.
- Möbius number :  $(n - 1)^{n-2}$  [McCammond and Meier 2004]

## Goal:

- New computation of the homology dimension
- Computation of the action of the symmetric group on the homology (Conjecture of Chapoton 2007)

## Theorem ([MM04])

*The poset  $\widehat{\text{HT}}_n$  is Cohen-Macaulay.*

## Theorem ([MM04])

*The poset  $\widehat{HT}_n$  is Cohen-Macaulay.*

## Corollary

*The character for the action of the symmetric group on  $\tilde{H}_{n-3}$  is given in terms of characters for the action of the symmetric group on  $C_k$  by:*

$$\chi_{\tilde{H}_{n-3}} = (-1)^{n-3} \sum_{k=-1}^{n-3} (-1)^k \chi_{C_k}, \text{ where } n = \#I.$$

## 1 The hypertree poset


## 2 Computation of the homology of the hypertree poset


- Species
- Counting strict chains using large chains
- Pointed hypertrees
- Relations between chains of hypertrees
- Dimension of the homology

## 3 From the hypertree poset to rooted trees

## Examples of species

- $\{(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1)\}$  (Species of lists Assoc on  $\{1, 2, 3\}$ )
- $\{\{1, 2, 3\}\}$  (Species of non-empty sets Comm)
- $\{\{1\}, \{2\}, \{3\}\}$  (Species of pointed sets Perm)


-  (Species of rooted trees PreLie)


-  (Species of cycles)

These sets are the image by species of the set  $\{1, 2, 3\}$ .

## Examples of species

- $\{(\heartsuit, \spadesuit, \clubsuit), (\heartsuit, \clubsuit, \spadesuit), (\spadesuit, \heartsuit, \clubsuit), (\spadesuit, \clubsuit, \heartsuit), (\clubsuit, \heartsuit, \spadesuit), (\clubsuit, \spadesuit, \heartsuit)\}$   
(Species of lists Assoc on  $\{\clubsuit, \heartsuit, \spadesuit\}$ )
- $\{\{\heartsuit, \spadesuit, \clubsuit\}\}$  (Species of non-empty sets Comm)
- $\{\{\heartsuit\}, \{\spadesuit\}, \{\clubsuit\}\}$  (Species of pointed sets Perm)

-  (Species of rooted trees PreLie)

-  (Species of cycles)

These sets are the image by species of the set  $\{\clubsuit, \heartsuit, \spadesuit\}$ .

# What are species?

## Definition

A *species*  $F$  is a functor from the category of finite sets and bijections to itself. To a finite set  $I$ , the species  $F$  associates a finite set  $F(I)$  independent from the nature of  $I$ .



# What are species?

## Definition

A *species*  $F$  is a functor from the category of finite sets and bijections to itself. To a finite set  $I$ , the species  $F$  associates a finite set  $F(I)$  independent from the nature of  $I$ .


Species = Construction plan, such that the set obtained is invariant by relabelling

# What are species?

## Definition

A *species*  $F$  is a functor from the category of finite sets and bijections to itself. To a finite set  $I$ , the species  $F$  associates a finite set  $F(I)$  independent from the nature of  $I$ .

Species = Construction plan, such that the set obtained is invariant by relabelling




# What are species?

## Definition

A *species*  $F$  is a functor from the category of finite sets and bijections to itself. To a finite set  $I$ , the species  $F$  associates a finite set  $F(I)$  independent from the nature of  $I$ .

Species = Construction plan, such that the set obtained is invariant by relabelling



# What are species?


## Definition

A *species*  $F$  is a functor from the category of finite sets and bijections to itself. To a finite set  $I$ , the species  $F$  associates a finite set  $F(I)$  independent from the nature of  $I$ .

## Counterexamples

The following sets are not obtained using species:

- $\{(1, \mathbf{3}, 2), (2, 1, \mathbf{3}), (2, \mathbf{3}, 1), (3, 1, \mathbf{2})\}$  (set of permutations of  $\{1, 2, 3\}$  with exactly 1 descent)
- (graph of divisibility of  $\{1, 2, 3, 4, 5, 6\}$ )



# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)


# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)

Example: Derivative of the species of cycles on  $I = \{\heartsuit, \spadesuit, \clubsuit\}$



# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)
- $(F + G)(I) = F(I) \sqcup G(I)$ , (addition)

# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)
- $(F + G)(I) = F(I) \sqcup G(I)$ , (addition)
- $(F \times G)(I) = \sum_{I_1 \sqcup I_2 = I} F(I_1) \times G(I_2)$ , (product)



# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)
- $(F + G)(I) = F(I) \sqcup G(I)$ , (addition)
- $(F \times G)(I) = \sum_{I_1 \sqcup I_2 = I} F(I_1) \times G(I_2)$ , (product)
- $(F \circ G)(I) = \bigsqcup_{\pi \in \mathcal{P}(I)} F(\pi) \times \prod_{J \in \pi} G(J)$ , (substitution) where  $\mathcal{P}(I)$  runs on the set of partitions of  $I$ .


# Operations on species and associated series

## Proposition

Let  $F$  and  $G$  be two species. The following operations can be defined on them:

- $F'(I) = F(I \sqcup \{\bullet\})$ , (derivative)
- $(F + G)(I) = F(I) \sqcup G(I)$ , (addition)
- $(F \times G)(I) = \sum_{I_1 \sqcup I_2 = I} F(I_1) \times G(I_2)$ , (product)
- $(F \circ G)(I) = \bigsqcup_{\pi \in \mathcal{P}(I)} F(\pi) \times \prod_{J \in \pi} G(J)$ , (substitution) where  $\mathcal{P}(I)$  runs on the set of partitions of  $I$ .

Example of substitution: Rooted trees of lists on  $I = \{1, 2, 3, 4\}$



## Definition

To a species  $F$ , we associate its *generating series*:

$$C_F(x) = \sum_{n \geq 0} \#F(\{1, \dots, n\}) \frac{x^n}{n!}.$$

## Examples of generating series:

- The generating series of the species of lists is  $C_{\text{Assoc}} = \frac{1}{1-x}$ .
- The generating series of the species of non-empty sets is  $C_{\text{Comm}} = \exp(x) - 1$ .
- The generating series of the species of pointed sets is  $C_{\text{Perm}} = x \cdot \exp(x)$ .
- The generating series of the species of rooted trees is  $C_{\text{PreLie}} = \sum_{n \geq 0} n^{n-1} \frac{x^n}{n!}$ .
- The generating series of the species of cycles is  $C_{\text{Cycles}} = -\ln(1-x)$ .

## Definition

The *cycle index series* of a species  $F$  is the formal power series in an infinite number of variables  $\mathfrak{p} = (p_1, p_2, p_3, \dots)$  defined by:

$$Z_F(\mathfrak{p}) = \sum_{n \geq 0} \frac{1}{n!} \left( \sum_{\sigma \in \mathfrak{S}_n} F^\sigma p_1^{\sigma_1} p_2^{\sigma_2} p_3^{\sigma_3} \dots \right),$$

- with  $F^\sigma$ , is the set of  $F$ -structures fixed under the action of  $\sigma$ ,
- and  $\sigma_i$ , the number of cycles of length  $i$  in the decomposition of  $\sigma$  into disjoint cycles.

## Examples

- The cycle index series of the species of lists is  $Z_{\text{Assoc}} = \frac{1}{1-p_1}$ .
- The cycle index series of the species of non empty sets is  $Z_{\text{Comm}} = \exp\left(\sum_{k \geq 1} \frac{p_k}{k}\right) - 1$ .

# Operations on cycle index series

Operations on species give operations on their cycle index series:

## Proposition

Let  $F$  and  $G$  be two species. Their cycle index series satisfy:

$$\begin{aligned} Z_{F+G} &= Z_F + Z_G, & Z_{F \times G} &= Z_F \times Z_G, \\ Z_{F \circ G} &= Z_F \circ Z_G, & Z_{F'} &= \frac{\partial Z_F}{\partial p_1}. \end{aligned}$$

## Definition

The *suspension*  $\Sigma$  of a cycle index series  $f(p_1, p_2, p_3, \dots)$  is defined by:

$$\Sigma f = -f(-p_1, -p_2, -p_3, \dots).$$

# Counting strict chains using large chains

Let  $I$  be a finite set of cardinality  $n$ .

## Definition

A *large  $k$ -chain* of hypertrees on  $I$  is a  $k$ -tuple  $(a_1, \dots, a_k)$ , where  $a_i$  are hypertrees on  $I$  and  $a_i \preceq a_{i+1}$ .

# Counting strict chains using large chains

Let  $I$  be a finite set of cardinality  $n$ .

## Definition

A *large  $k$ -chain* of hypertrees on  $I$  is a  $k$ -tuple  $(a_1, \dots, a_k)$ , where  $a_i$  are hypertrees on  $I$  and  $a_i \preceq a_{i+1}$ .

Let  $M_{k,s}$  be the set of words on  $\{0, 1\}$  of length  $k$ , with  $s$  letters "1". The species  $\mathcal{M}_{k,s}$  is defined by:

$$\begin{cases} \emptyset & \mapsto M_{k,s}, \\ V \neq \emptyset & \mapsto \emptyset. \end{cases}$$

# Counting strict chains using large chains

Let  $I$  be a finite set of cardinality  $n$ .

## Definition

A *large  $k$ -chain* of hypertrees on  $I$  is a  $k$ -tuple  $(a_1, \dots, a_k)$ , where  $a_i$  are hypertrees on  $I$  and  $a_i \preceq a_{i+1}$ .

Let  $M_{k,s}$  be the set of words on  $\{0, 1\}$  of length  $k$ , with  $s$  letters "1". The species  $\mathcal{M}_{k,s}$  is defined by:

$$\begin{cases} \emptyset & \mapsto M_{k,s}, \\ V \neq \emptyset & \mapsto \emptyset. \end{cases}$$

## Proposition

The species  $\mathcal{H}_k$  of large  $k$ -chains and  $\mathcal{HS}_i$  of strict  $i$ -chains are related by:

$$\mathcal{H}_k \cong \sum_{i \geq 0} \mathcal{HS}_i \times \mathcal{M}_{k,i}.$$



## Proposition

The species  $\mathcal{H}_k$  of large  $k$ -chains and  $\mathcal{HS}_i$  of strict  $i$ -chains are related by:

$$\mathcal{H}_k \cong \sum_{i \geq 0} \mathcal{HS}_i \times \mathcal{M}_{k,i}.$$

Proof.

Deletion of repetitions

$$(a_1, \dots, a_k)$$

$$(a_{j_1}, \dots, a_{j_i})$$

$u_j = 0$  if  $a_j = a_{j-1}$ , 1 otherwise

$$(u_1, \dots, u_k)$$

with  $a_0 = \hat{0}$ . □

The previous proposition gives, for all integer  $k > 0$ :

$$\chi_k = \sum_{i=0}^{n-2} \binom{k}{i} \chi_i^s.$$

The previous proposition gives, for all integer  $k > 0$ :

$$\chi_k = \sum_{i=0}^{n-2} \binom{k}{i} \chi_i^s.$$

$\chi_k$  is a polynomial  $P(k)$  in  $k$  which gives, once evaluated in  $-1$ , the character:

### Corollary

$$\chi_{\tilde{H}_{n-3}} = (-1)^n P(-1) =: (-1)^n \chi_{-1}$$

The hypertrees will now be on  $n$  vertices.


# Pointed hypertrees

## Definition

Let  $H$  be a hypertree on  $I$ .  $H$  is:

- *rooted* in a vertex  $s$  if the vertex  $s$  of  $H$  is distinguished,
- *edge-pointed* in an edge  $a$  if the edge  $a$  of  $H$  is distinguished,
- *rooted edge-pointed* in a vertex  $s$  in an edge  $a$  if the edge  $a$  of  $H$  and a vertex  $s$  of  $a$  are distinguished.

## Example of pointed hypertrees



## Proposition: Dissymmetry principle

The species of hypertrees and of rooted hypertrees are related by:

$$\mathcal{H} + \mathcal{H}^{pa} = \mathcal{H}^p + \mathcal{H}^a.$$

We write:

- $\mathcal{H}_k$ , the species of large  $k$ -chains of hypertrees,
- $\mathcal{H}_k^{pa}$ , the species of large  $k$ -chains of hypertrees whose minimum is rooted edge-pointed,
- $\mathcal{H}_k^p$ , the species of large  $k$ -chains of hypertrees whose minimum is rooted,
- $\mathcal{H}_k^a$ , the species of large  $k$ -chains of hypertrees whose minimum is edge-pointed.

## Corollary ([Oge13])

*The species of large  $k$ -chains of hypertrees are related by:*


$$\mathcal{H}_k + \mathcal{H}_k^{pa} = \mathcal{H}_k^p + \mathcal{H}_k^a.$$

## Last but not least type of hypertrees

### Definition

A *hollow hypertree* on  $n$  vertices ( $n \geq 2$ ) is a hypertree on the set  $\{\#, 1, \dots, n\}$ , such that the vertex labelled by  $\#$ , called the *gap*, belongs to one and only one edge.

### Example of a hollow hypertree



We denote by  $\mathcal{H}_k^c$ , the species of large  $k$ -chains of hypertrees whose minimum is a hollow hypertree.

# Relations between species of hypertrees

## Theorem

The species  $\mathcal{H}_k$ ,  $\mathcal{H}_k^P$  and  $\mathcal{H}_k^C$  satisfy:

$$\mathcal{H}_k^P = X \times \mathcal{H}'_k \quad (1)$$

$$\mathcal{H}_k^P = X \times \text{Comm} \circ \mathcal{H}_k^C + X, \quad (2)$$

$$\mathcal{H}_k^C = \text{Comm} \circ \mathcal{H}_{k-1}^C \circ \mathcal{H}_k^P, \quad (3)$$

$$\mathcal{H}_k^a = (\mathcal{H}_{k-1} - x) \circ \mathcal{H}_k^P, \quad (4)$$

$$\mathcal{H}_k^{pa} = (\mathcal{H}_{k-1}^P - x) \circ \mathcal{H}_k^P. \quad (5)$$

# Relations between species of hypertrees

## Theorem

The species  $\mathcal{H}_k$ ,  $\mathcal{H}_k^p$  and  $\mathcal{H}_k^c$  satisfy:

$$\mathcal{H}_k^p = X \times \mathcal{H}'_k \quad (1)$$

$$\mathcal{H}_k^p = X \times \text{Comm} \circ \mathcal{H}_k^c + X, \quad (2)$$

$$\mathcal{H}_k^c = \text{Comm} \circ \mathcal{H}_{k-1}^c \circ \mathcal{H}_k^p, \quad (3)$$

$$\mathcal{H}_k^a = (\mathcal{H}_{k-1} - x) \circ \mathcal{H}_k^p, \quad (4)$$

$$\mathcal{H}_k^{pa} = (\mathcal{H}_{k-1}^p - x) \circ \mathcal{H}_k^p. \quad (5)$$

## Proof.


- 1 Rooting a species  $F$  is the same as multiplying the singleton species  $X$  by the derivative of  $F$ ,



## Second part of the proof.

We separate the root and every edge containing it, putting gaps where the root was,


$$\mathcal{H}_k^p = X \times \text{Comm} \circ \mathcal{H}_k^c + X,$$



3 Hollow case:

$$\mathcal{H}_k^c = \mathcal{H}_k^{cm} \circ \mathcal{H}_k^p, \quad (6)$$

$$\mathcal{H}_k^{cm} = \text{Comm} \circ \mathcal{H}_{k-1}^c. \quad (7)$$



# Dimension of the homology

## Proposition

The generating series of the species  $\mathcal{H}_k$ ,  $\mathcal{H}_k^p$  and  $\mathcal{H}_k^c$  satisfy:

$$\mathcal{C}_k^p = x \cdot \exp \left( \frac{\mathcal{C}_{k-1}^p \circ \mathcal{C}_k^p}{\mathcal{C}_k^p} - 1 \right), \quad (8)$$

$$\mathcal{C}_k^a = (\mathcal{C}_{k-1} - x)(\mathcal{C}_k^p), \quad (9)$$

$$\mathcal{C}_k^{pa} = (\mathcal{C}_{k-1}^p - x)(\mathcal{C}_k^p), \quad (10)$$

$$x \cdot \mathcal{C}_k' = \mathcal{C}_k^p, \quad (11)$$

$$\mathcal{C}_k + \mathcal{C}_k^{pa} = \mathcal{C}_k^p + \mathcal{C}_k^a. \quad (12)$$

## Lemma

The generating series of  $\mathcal{H}_0$  and  $\mathcal{H}_0^P$  are given by:

$$C_0 = \sum_{n \geq 1} \frac{x^n}{n!} = e^x - 1,$$

$$C_0^P = xe^x.$$

## Lemma

The generating series of  $\mathcal{H}_0$  and  $\mathcal{H}_0^P$  are given by:

$$C_0 = \sum_{n \geq 1} \frac{x^n}{n!} = e^x - 1,$$

$$C_0^P = xe^x.$$

This implies with the previous theorem:

## Theorem ([MM04])

*The dimension of the only homology group of the hypertree poset is  $(n - 1)^{n-2}$ .*

This dimension is the dimension of the vector space  $\text{PreLie}(n-1)$  whose basis is the set of rooted trees on  $n - 1$  vertices.

- 1 The hypertree poset
- 2 Computation of the homology of the hypertree poset
- 3 From the hypertree poset to rooted trees
  - PreLie species
  - Character for the action of the symmetric group on the homology of the poset

# From the hypertree poset to rooted trees

- 1 This dimension is the dimension of the vector space  $\text{PreLie}(n-1)$  whose basis is the set of rooted trees on  $n - 1$  vertices.  
The operad (a species with more properties on substitution) whose vector space are  $\text{PreLie}(n)$  is  $\text{PreLie}$ .
- 2 This operad is **anticyclic** ([Cha05]): There is an action of the symmetric group  $\mathfrak{S}_n$  on  $\text{PreLie}(n - 1)$  which extends the one of  $\mathfrak{S}_{n-1}$ .
- 3 Moreover, there is an **action** of  $\mathfrak{S}_n$  on the homology of the poset of hypertrees on  $n$  vertices.

## Question

Is the action of  $\mathfrak{S}_n$  on  $\text{PreLie}(n-1)$  the same as the action on the homology of the poset of hypertrees on  $n$  vertices?

# Character for the action of the symmetric group on the homology of the poset

Using relations on species established previously, we obtain:

## Proposition

The series  $Z_k$ ,  $Z_k^p$ ,  $Z_k^a$  and  $Z_k^{pa}$  satisfy the following relations:

$$Z_k + Z_k^{pa} = Z_k^p + Z_k^a, \quad (13)$$

$$Z_k^p = p_1 + p_1 \times \text{Comm} \circ \left( \frac{Z_{k-1}^p \circ Z_k^p - Z_k^p}{Z_k^p} \right), \quad (14)$$

$$Z_k^a + Z_k^p = Z_{k-1} \circ Z_k^p, \quad (15)$$

$$Z_k^{pa} + Z_k^p = Z_{k-1}^p \circ Z_k^p, \text{ and } p_1 \frac{\partial Z_k}{\partial p_1} = Z_k^p. \quad (16)$$



## Theorem ([Oge13], conjecture of [Cha07])

The cycle index series  $Z_{-1}$ , which gives the character for the action of  $\mathfrak{S}_n$  on  $\tilde{H}_{n-3}$ , is linked with the cycle index series  $M$  associated with the anticyclic structure of PreLie by:

$$Z_{-1} = p_1 - \Sigma M = \text{Comm} \circ \Sigma \text{PreLie} + p_1 (\Sigma \text{PreLie} + 1). \quad (17)$$

The cycle index series  $Z_{-1}^P$  is given by:

$$Z_{-1}^P = p_1 (\Sigma \text{PreLie} + 1). \quad (18)$$

## Theorem ([Oge13], conjecture of [Cha07])

The cycle index series  $Z_{-1}$ , which gives the character for the action of  $\mathfrak{S}_n$  on  $\tilde{H}_{n-3}$ , is linked with the cycle index series  $M$  associated with the anticyclic structure of PreLie by:

$$Z_{-1} = p_1 - \Sigma M = \text{Comm} \circ \Sigma \text{PreLie} + p_1 (\Sigma \text{PreLie} + 1). \quad (17)$$

The cycle index series  $Z_{-1}^P$  is given by:

$$Z_{-1}^P = p_1 (\Sigma \text{PreLie} + 1). \quad (18)$$

## Proof.

Sketch of the proof

- 1 Computation of  $Z_0 = \text{Comm}$  and  $Z_0^P = \text{Perm} = p_1 + p_1 \times \text{Comm}$
- 2 Replaced in the formula giving  $Z_0^P$  in terms of itself and  $Z_{-1}^P$

$$Z_0^P = p_1 + p_1 \times \text{Comm} \circ \left( \frac{Z_{-1}^P \circ Z_0^P - Z_0^P}{Z_0^P} \right),$$

## Second part of the proof.

- ③ As  $\Sigma \text{PreLie} \circ \text{Perm} = \text{Perm} \circ \Sigma \text{PreLie} = p_1$ , according to [Cha07], we get:

$$Z_{-1}^P = p_1 (\Sigma \text{PreLie} + 1).$$

- ④ The dissymetry principle associated with the expressions gives:

$$\text{Comm} + Z_{-1}^P \circ \text{Perm} - \text{Perm} = \text{Perm} + Z_{-1} \circ \text{Perm} - \text{Perm}.$$

- ⑤ Thanks to equation [Cha05, equation 50], we conclude:

$$\Sigma M - 1 = -p_1 \left( -1 + \Sigma \text{PreLie} + \frac{1}{\Sigma \text{PreLie}} \right).$$



# Open questions

A remplir!!!!

Thank you for your attention !

[Oge13] [Bérénice Oger](#) Action of the symmetric groups on the homology of the hypertree posets. [Journal of Algebraic Combinatorics](#), february 2013.

# Bibliography I



Claude Berge :

Hypergraphs, volume 45 de North-Holland Mathematical Library.  
North-Holland Publishing Co., Amsterdam, 1989.  
Combinatorics of finite sets.



Frédéric Chapoton :

On some anticyclic operads.

Algebr. Geom. Topol., 5:53–69 (electronic), 2005.



Frédéric Chapoton :

Hyperarbres, arbres enracinés et partitions pointées.

Homology, Homotopy Appl., 9(1):193–212, 2007.

<http://www.intlpress.com/hha/v9/n1/>.

# Bibliography II



Jon McCammond et John Meier :

The hypertree poset and the  $l^2$ -Betti numbers of the motion group of the trivial link.

[Math. Ann.](#), 328(4):633–652, 2004.



Bérénice Oger :

Action of the symmetric groups on the homology of the hypertree posets.

[J. Algebraic Combin.](#), 38(4):915–945, 2013.

# Eccentricity

## Definition

The *eccentricity* of a vertex or an edge is the maximal number of vertices on a walk without repetition to another vertex.

The *center* of a hypertree is the vertex or the edge with minimal eccentricity.

## Example of eccentricity

